## CLAIMS

We claim:

- 1. An isolated polynucleotide encoding a ligand-binding receptor polypeptide, said polypeptide being defined by amino acid residues 18 to 228 of SEQ ID NO:2.
- 2. An isolated polynucleotide according to claim 1 wherein said polypeptide further comprises a transmembrane domain.
- 3. An isolated polynucleotide according to claim 2 wherein said transmembrane domain comprises residues 229 to 251 of SEQ ID NO:2.
- 4. An isolated polynucleotide according to claim 2 wherein said polypeptide further comprises an intracellular domain.
- 5. An isolated polynucleotide according to claim 4 wherein said intracellular domain comprises residues 252 to 574 of SEQ ID NO:2.
- 6. An isolated polynucleotide according to claim 1 which is a DNA as shown in SEQ ID NO:1 from nucleotide 34 to nucleotide 1755.
- 7. An isolated polynucleotide according to claim 1 wherein said polypeptide further comprises an affinity tag.
- 8. An isolated polynucleotide according to claim 7 wherein said affinity tag is polyhistidine, protein A, glutathione S transferase, substance P, or an immunoglobulin heavy chain constant region.



- 9. An isolated polynucleotide according to claim 1 wherein said polynucleotide is DNA.
- An isolated polynucleotide encoding a polypeptide selected from a group defined SEQ ID NO:2 consisting of residues 1 to 228, residues 1 to 251, residues 1 to 574, residues 2 to 228, residues 2 to 251, residues 2 to 574, residues 229 to 251, residues 229 to 574 and residues 252 to 574.
- 11. An expression vector comprising the following operably linked elements:
  - a transcription promoter;
- a DNA segment encoding a ligand-binding receptor polypeptide, said polypeptide being defined by amino residues 18 to 228 of SEQ ID NO:2; and
  - a transcription terminator.
- 12. An expression vector according to claim 11 wherein said polypeptide further comprises a signal sequence.
- 13. An expression vector according to claim 11 wherein said polypeptide further comprises a transmembrane domain.
- 14. An expression vector according to claim 11 wherein said transmembrane domain comprises residues 229 to 251 of SEQ ID NO:2.
- 15. An expression vector according to claim 13 wherein said polypeptide further comprises an intracellular domain.
- 16. An expression vector according to claim 15 wherein said intracellular domain comprises residues 252 to 574 of SEQ ID NO:2.

- An expression vector according to claim 11 wherein further comprising a DNA sequence encoding an affinity tag.
- 18. An expression vector according to claim 17 wherein the affinity tag is an immunoglobulin  $F_{\rm C}$  polypeptide.
- 19. A transformed or transfected cell into which has been introduced an expression vector according to claim 11, wherein said cell expresses a receptor polypeptide encoded by the DNA segment.
- 20. An isolated polypertide defined by residues 18-228 of SEQ ID NO: 2.
- 21. The isolated polypeptide of claim 20 further containing either a sequence which defines a transmembrane domain or a sequence which defines an intracellular domain or both.
- 22. The isolated polypeptide of claim 23 wherein the transmembrane domain is defined by amino acid residues 229-251 of SEQ ID NO: 2 and the intracellular domain is defined by amino acid residues 252-574 of SEQ ID NO:2.
- 23. An isolated polypeptide according to claim 20 further containing a sequence which defines an affinity tag.
- 24. A method for detecting a ligand within a test sample, comprising contacting a test sample with a polypeptide comprising residues 18 to 228 of SEQ ID NO:2; and detecting binding of said polypeptide to a ligand in the sample.

5,0

- 25. An antibody that specifically binds to a polypeptide of claim 20.
- 26. An anti-idiotypic antibody which binds to an antigenic binding site of an antibody of claim 25.
- 27. An isolated polypeptide selected from the group consisting of residues 1 to 228, residues 1 to 251, residues 1 to 574 residues 2 to 228, residues 2 to 551, and residues 2 to 574 of SEQ ID NO: 2.

add of